

۲. عالی (خوبه) ضربه سیف کندی مقبره

Date

No

نام خدا

سارنا ایبری شهبازی

الف) $\tan \frac{\pi}{6} + \sin \frac{\pi}{6} \cos \frac{\pi}{6} \rightarrow -1 + (-\frac{\sqrt{2}}{2} \times \frac{\sqrt{2}}{2}) = -\frac{1}{2}$ ①

ب) $\tan \frac{\pi}{6} \sin \frac{\pi}{6} + \cos \frac{\pi}{6} \rightarrow (\frac{\sqrt{2}}{2} \times \frac{\sqrt{2}}{2}) - \frac{1}{2} = \frac{1}{2}$ ②

~~$\cos x = \frac{a}{r}$~~ ~~$\sin x = \frac{b}{r}$~~ ~~$\tan x = \frac{b}{a}$~~ ~~$\cot x = \frac{a}{b}$~~ ~~$\sec x = \frac{r}{a}$~~ ~~$\csc x = \frac{r}{b}$~~


$\cos x = \frac{a}{r}$ / $\sin x = \frac{b}{r}$

$\frac{\cos x}{\sin x} = \frac{a}{b}$ ③

$\sin x = r \cos x$ $\tan x = \frac{b}{a}$

cos
sin

$\cos = \frac{a}{r}$



$\cos x = -\frac{1}{\sqrt{2}} = -\frac{\sqrt{2}}{2}$ ④

الف) $\sin x = \frac{\sqrt{2}}{1}$
 $\cos x = \frac{\sqrt{2}}{1}$

$\cos \frac{\pi}{6} \cos x - \sin \frac{\pi}{6} \sin x \rightarrow \frac{1}{1} - \frac{1}{1} = \frac{1}{1} = \frac{1}{1}$ ⑤

~~$\tan x = \frac{b}{a}$~~ ~~$\cot x = \frac{a}{b}$~~ ~~$\sec x = \frac{r}{a}$~~ ~~$\csc x = \frac{r}{b}$~~

ب) $\tan x = \frac{1}{\sqrt{2}}$
 $\sin x = \frac{1}{\sqrt{2}} = \frac{\sqrt{2}}{2}$, $\cos x = \frac{\sqrt{2}}{2}$

$\sin \frac{\pi}{6} \cos x + \sin x \cos \frac{\pi}{6} \rightarrow (\frac{\sqrt{2}}{2}) (\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}) = \frac{1}{1} = \frac{1}{1}$ ⑥

$$\sin^2 x + \sin^2 x + \cos^2 x = \frac{r}{r} \rightarrow \sin x = \frac{1}{\sqrt{r}} \quad (2)$$

$$\cos^2 x = \frac{r}{r} \rightarrow \cos x = \frac{\sqrt{r}}{\sqrt{r}}$$


$$\tan^2 x = \left(\frac{\frac{1}{\sqrt{r}}}{\frac{\sqrt{r}}{\sqrt{r}}} \right) = \frac{1}{r}$$



$$\frac{1}{r} \times \sqrt{r} \times r \times \sin \alpha = \frac{r}{r} \quad (4)$$

$$\sin \alpha = \frac{1}{r} \times \frac{1}{\sqrt{r}} = \frac{1}{r\sqrt{r}} = \frac{\sqrt{r}}{r} \rightarrow \alpha = \gamma^\circ$$

$$\alpha = 12^\circ$$



$$S_{ABCD} = r \left(\frac{1}{r} \times r \times r \times \sin 12^\circ \right) = \frac{r}{r} \quad (5)$$

$$P_{ABCD} = r \left(\frac{1}{r} \times r \times r \right) = \frac{r}{r} \quad (6)$$

$$S_{ABC} - S_{ADE} = \frac{1}{r} \times \frac{r}{r}$$

$$\left(\frac{1}{r} \times r \times r \times \sin A \right) - \left(\frac{1}{r} \times r \times r \times \sin A \right) = \frac{1}{r} \times \frac{r}{r}$$

$$\sin A (r - r) = \frac{1}{r} \times \frac{r}{r} \rightarrow \sin A = \frac{1}{r} \rightarrow A = \gamma^\circ$$

$$\tan \gamma^\circ = \frac{\frac{1}{\sqrt{r}}}{\frac{\sqrt{r}}{\sqrt{r}}}$$

$$\frac{S_{BMN}}{S_{ABC}} = \frac{1}{r} \rightarrow \frac{\frac{1}{r} \times BM \times BN \times \sin B}{\frac{1}{r} \times AB \times BC \times \sin B} = \frac{1}{r}$$

$$\frac{BM}{AB} = \frac{1}{r}$$

$$\frac{BM}{AM} = \frac{BM}{(AB - BM)} = \frac{1}{r}$$

$$\frac{BM}{AB} \times \frac{BN}{BC} = \frac{1}{r}$$

$$\frac{|\sin \alpha|}{\cos \alpha} = -\frac{1}{\cos \alpha} = -\frac{1}{\cos \alpha} = -\frac{\sin \alpha}{\cos \alpha} \rightarrow \sin \alpha < 0 \quad (1)$$

$$\frac{1}{\cos \alpha} = \frac{\sin \alpha}{\cos \alpha} = \frac{1 + \sin \alpha}{\cos \alpha} \quad (2)$$