

$$y = \sqrt{x f(x)} \quad x f(x) \geq 0 \quad \begin{array}{c} -a \quad 0 \quad r \quad a \\ \hline \overline{0} + \overline{0} + \overline{0} + \overline{0} \end{array} \quad (5)$$

$$D_f = [r, a] \cup [-a, 0]$$

$$y = \sqrt{\frac{-x}{f(x)}} \quad \frac{-x}{f(x)} \geq 0 \quad \begin{array}{c} -a \quad -r \quad 0 \quad r \quad a \\ \hline \overline{-} + \overline{+} + \overline{+} + \overline{-} \end{array} \quad (1, a)$$

$$D_f = (-r, r) - \{0\} \rightarrow \text{عنصر صفر}$$

$$f(x) - f(r) = x^2 - rx + \epsilon \xrightarrow{x=r} -f(r) = -r \rightarrow f(r) = -r \leq r$$

$$\xrightarrow{x=-r} f(-r) = \epsilon + 4 = 10$$

$$\begin{cases} x - \sqrt{x} + \epsilon & : x > r \\ rx + r & , x \leq r \end{cases} \quad f(f(a)) + f(f(1)) \quad (5)$$

$$f(a) = a - \sqrt{a} = r \xrightarrow{f(r)} f(r) = v \quad f(1) = a \xrightarrow{f(a)} f(a) = r$$

$$v + r = 9$$

$$f(x-1) = ax^2 + a - 2ax - bx - b + r \quad (5)$$

$$-f(x) + f(x-1) = ax^2 + a - 2ax - bx - b + r - ax^2 + bx - r$$

$$\rightarrow a - 2ax - b = 4x + r \rightarrow -2ax = 4x \Rightarrow a = -r$$

$$\Rightarrow b = -a \rightarrow -r - a = -1$$

Hashtag

H A S H T A G

$$f(x) = \frac{x^5 + 5x + 4}{x^5 + 5x + 4} = \frac{(x+1)^5 + 1}{(x+1)^5 + 1} \xrightarrow{f(1-1)} \quad \textcircled{5} -4$$

$$\frac{(\sqrt{5})^5 + 1}{(\sqrt{5})^5 + 4} = \frac{6}{4} = \frac{3}{2}$$

$$f\left(x - \frac{1}{x}\right) = \frac{x^5 + 1}{x^5} = x^5 + \frac{1}{x^5} \xrightarrow{f\left(x - \frac{1}{x}\right) = \frac{\left(x - \frac{1}{x}\right)^5 + 1}{\left(x - \frac{1}{x}\right)^5 + 4} = V$$

$$\hookrightarrow \left(x - \frac{1}{x}\right)^5 = x^5 + \frac{1}{x^5} - 4$$

$$\Rightarrow f(-2) = 9 + 2 = 11$$

$$g(x) = \sqrt{9 - x^2} \quad 9 - x^2 \geq 0 \quad x^2 \leq 9 \rightarrow -3 \leq x \leq 3 \quad \textcircled{5} -1$$

$$\text{الف) } \frac{f}{g} = \left\{ \left(0, \frac{2}{3}\right), \left(1, \frac{-1}{\sqrt{2}}\right), \left(2, \frac{0}{\sqrt{5}}\right) \right\}$$

$$\text{ب) } \frac{g}{f} = \left\{ \left(0, \frac{3}{2}\right), \left(1, -\frac{\sqrt{2}}{1}\right), \left(2, \frac{\sqrt{5}}{0}\right) \right\}$$

نقطة

$$\text{الف) } \{(2, 2), (3, 1), (-5, 4), (1, -1)\} \quad \textcircled{5} -4$$

$$\text{ب) } \{(2, 2), (3, 5), (-5, 3), (1, -1)\}$$

$$\text{ج) } \{(2, 4), (3, 9), (-5, 13), (1, 13)\}$$

$$\text{د) } \left\{ \left(1, 1\right), \left(\frac{5}{2}, 4\right), \left(-\frac{5}{2}, 2\right), \left(\frac{1}{2}, -2\right) \right\}$$

Subject:

Date: / /

$$g(x) = \{ \widetilde{(1, 0)}, \widetilde{(2, 1)}, \widetilde{(-1, 4)}, \widetilde{(-2, -3)}, \widetilde{(3, -1)} \} \quad \text{--- 10} \quad \textcircled{P}$$

$$f(x) = \{ \widetilde{(2, 3)}, \widetilde{(4, 2)}, \widetilde{(1, 4)}, \widetilde{(3, 1)}, \widetilde{(2, 2)} \}$$

$$\text{الف) } f - g = \{ (1, 4), (2, 2), (3, 2) \}$$

$$\text{ب) } \frac{f}{g} = \{ \underbrace{(1, \frac{4}{0})}_{\text{غير معرف}}, (2, 4), (3, \frac{2}{-1}) \}$$