

نام و نام خانوادگی: ... پسر ... پاسخنانه تشریحی تکلیف شماره ۱ ... کلاس: ...

الف $ext = \begin{bmatrix} -3 \\ -14 \end{bmatrix}$ $x = \frac{-b}{2a} = \frac{-4}{2} = -2$ $y = x^2 + 4x + 5$

$y = (-3)^2 + 4(-3) + 5 = -14$ جاگذاری



نمودار طول از مبدأهای ۵- و ۱- دارد


ب) $a > 0 \Rightarrow$ مینیمم دارد

اگر $x = 0 \Rightarrow y = 0^2 + 4(0) + 5 = 5$ عرض از مبدأ ۵ دارد

$x = -5$ و $x = 1$
 $y = 0 \Rightarrow x^2 + 4x + 5 = 0$
 $0 = (x+5)(x+1) = 0$

عرض از مبدأ آن هم -3 است $y = -x^2 + 4x - 3$

الف $ext \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ $x = \frac{-b}{2a} = \frac{-4}{-2} = 2$ $y = -(2)^2 + 4(2) - 3 = 1$ جاگذاری



است max $a < 0$

ب) $y = 0 \Rightarrow -x^2 + 4x - 3 = 0 \Rightarrow x^2 - 4x + 3 = 0$
 $x^2 - 4x + 3 = 0$
 $a + c + b = 0$
 $\Rightarrow x = 1$ و $x = 3$

الف $\Rightarrow \Delta > 0 \Rightarrow b^2 - 4ac > 0 \Rightarrow (-3)^2 - 4(2)(1) > 0$
 $\Rightarrow 9 - 8 > 0 \Rightarrow 1 > 0$

ب) $\Delta = 0 \Rightarrow b^2 - 4ac = 0$
 $\Rightarrow (-3)^2 - 4(2)(1) = 0$
 $\Rightarrow 9 - 8 = 1 \neq 0$

2) $\Delta < 0 \Rightarrow b^2 - 4ac < 0 \Rightarrow (-3)^2 - 4(2)(1) < 0$
 $\Rightarrow 9 - 8 < 0$

3) $\Delta > 0 \Rightarrow b^2 - 4ac > 0 \Rightarrow (-3)^2 - 4(2)(1) > 0$
 $\Rightarrow 9 - 8 > 0$

الف $x^2 - 2x - 1 = 0 \Rightarrow x^2 - 2x + 1 = 1 \Rightarrow (x-1)^2 = 1$

$x = \frac{2 \pm \sqrt{4 + 4}}{2} = \frac{2 \pm \sqrt{8}}{2} = \frac{2 \pm 2\sqrt{2}}{2} = 1 \pm \sqrt{2}$

$\Rightarrow x - 1 = \pm \sqrt{1}$
 $\Rightarrow x = 1 + \sqrt{1}$ و $x = 1 - \sqrt{1}$
 $\Rightarrow x = 2$ و $x = 0$

ب) $x^2 - x - 1 = 0 \Rightarrow x^2 - x + 1/4 - 1 = 1/4$

$\Rightarrow x^2 - x + 1/4 = 1/4 + 1$
 $(x - 1/2)^2 = 5/4$
 $\Rightarrow x - 1/2 = \pm \sqrt{5/4}$
 $\Rightarrow x = 1/2 \pm \sqrt{5}/2$

الف $x^2 - 2x - 1 = 0$ $\Delta = b^2 - 4ac$
 $x = \frac{-b \pm \sqrt{\Delta}}{2a} = \frac{2 \pm \sqrt{4 + 4}}{2} = \frac{2 \pm \sqrt{8}}{2} = \frac{2 \pm 2\sqrt{2}}{2} = 1 \pm \sqrt{2}$

ب) $x^2 + x - 4 = 0$
 $x = \frac{-b \pm \sqrt{\Delta}}{2a} = \frac{-1 \pm \sqrt{1 + 16}}{2} = \frac{-1 \pm \sqrt{17}}{2}$

$\Rightarrow \begin{cases} x = \frac{-1 + \sqrt{17}}{2} \\ x = \frac{-1 - \sqrt{17}}{2} \end{cases}$

الف) $x^2 - 11x + 18 = 0 \Rightarrow (x-2)(x-9) = 0 \Rightarrow x=2 \vee x=9$

ب) $x^2 + 13x - 18 = 0 \Rightarrow (x+18)(x-1) = 0 \Rightarrow x=-18 \vee x=1$

٢

الف) $\Delta x^2 - 12x + 7 = 0 \Rightarrow x^2 - 12x + 36 = 0 \Rightarrow (x-6)(x-6) = 0$
 $\Rightarrow x=6$
 $\Rightarrow \begin{cases} x=6 \\ x=6 \end{cases} \Rightarrow \begin{cases} x=\frac{6}{6}=1 \\ x=\frac{6}{6}=1 \end{cases}$

٢

٧

ب) $\Delta x^2 - 10x + 1 = 0 \Rightarrow x^2 - 10x + 1 = 0 \Rightarrow (x-5)(x-5) = 0$
 $\Rightarrow \begin{cases} x=5 \\ x=5 \end{cases} \Rightarrow \begin{cases} x=\frac{5}{5}=1 \\ x=\frac{5}{5}=1 \end{cases}$

الف) $2x^2 - 5x + 2 = 0 \Rightarrow a+b+c=0 \Rightarrow \begin{cases} x=\frac{c}{a} = \frac{2}{2} \\ x=1 \end{cases}$

ب) $2x^2 + 5x + 2 = 0 \Rightarrow a+c=b \Rightarrow \begin{cases} x=-1 \\ x=-\frac{c}{a} = -\frac{2}{2} = -1 \end{cases}$

٢

٨

٢) $2x^2 - 5x + 1 = 0 \Rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{5 \pm \sqrt{25 - 4}}{4} = \frac{5 \pm \sqrt{21}}{4}$

$\Delta = b^2 - 4ac = 25 - 4 = 21 > 0$

$x^2 - 13x - 1 = 0 \Rightarrow \Delta = 169 + 4 = 173 > 0 \Rightarrow S = \frac{-b}{a} = 13, P = \frac{c}{a} = -1$

الف) $S^2 - P = 169 - (-1) = 170$

٢

٩

ب) $S^2 - 13S - 1 = 170 - 13(13) - 1 = 170 - 169 - 1 = 0$

الف) $\begin{pmatrix} V \\ \omega \end{pmatrix} + \begin{pmatrix} 9 \\ \mu \end{pmatrix} = \begin{pmatrix} V \\ \mu \end{pmatrix} + \begin{pmatrix} 9 \\ \mu \end{pmatrix} = \frac{V \times \mu}{\mu} + \frac{9 \times \mu}{\mu} = V + 9 = 10$

ب) $\begin{pmatrix} 9 \\ \mu \end{pmatrix} - \begin{pmatrix} 1 \\ \mu \end{pmatrix} = \frac{9 \times \mu}{\mu} - \frac{1 \times \mu}{\mu} = 8$

٢

١٠