

$$-c\varepsilon, \Delta + \pi\gamma = 1, \Delta$$

$$\textcircled{10}$$

$$\textcircled{1, \sqrt{0}}$$

$$f(1, \Delta) = \frac{1}{r} (1, \Delta) + 1 = \frac{\sqrt{0}}{2}$$

$$f(x) \begin{cases} -\frac{1}{r}x + 1 & ; -\varepsilon \leq x \leq r \\ rx + r & ; -1 \leq x \leq 0 \end{cases} \rightarrow f(1, \Delta) = -\frac{1}{r}(\frac{r}{r}) + 1 = \frac{1}{r}$$

$$f(x) = \frac{b}{r} \sin r\alpha + a$$

$$\textcircled{1}$$

$$f(x) = \frac{b}{r} \sin r\alpha + a$$

$$T = \frac{\pi}{r} - \frac{\pi}{10} = \frac{\pi}{10} = \frac{\pi}{10}$$

$$\frac{r\pi}{|r|} = \frac{r\pi}{0} = \frac{\pi}{0}$$

$$a = \frac{\varepsilon + 0}{r} = \gamma$$

$$|b| + r = \varepsilon = 1 \Rightarrow b = r$$

$$\begin{cases} a + \frac{b}{r} = r \\ a - \frac{b}{r} = \varepsilon \end{cases} \rightarrow a = r, b = 1 \rightarrow \frac{b}{a} = \frac{1}{r} = \textcircled{d}$$

$$\textcircled{9}$$

$$\frac{r\pi}{|b|} = \pi = \frac{\pi}{r} \Rightarrow b = r$$

$$\frac{r - \varepsilon}{r} = -1 = c$$

$$|a| - 1 = r = \varepsilon \Rightarrow a = r$$

$$-r \cos r\alpha - 1 = 0$$

$$\cos r\alpha = -\frac{1}{r}$$

$$1 + \tan r\alpha = \frac{1}{\cos r\alpha} = \frac{1}{-\frac{1}{r}} = -r$$

$$\Rightarrow |\tan r\alpha| = \sqrt{1}$$

$$r = AC + CB + \dots$$

$$\sqrt{1+1-r \cos \frac{\pi}{7}} + \sqrt{r-r \cos \frac{\pi}{7}} + \sqrt{r-r \cos \frac{\pi}{7}}$$

$$= \sqrt{r+\sqrt{r}} + \sqrt{r} + \sqrt{r}$$