

یازدهم ریاضی A

$$n \in \mathbb{N} \left\{ \begin{array}{l} a^n - r^n \\ (a-r)^n \end{array} \right\} \Rightarrow a^n - r^n > (a-r)^n \Rightarrow a^n - (a-r)^n > 0 \Rightarrow (a-r)(a^{n-1} + a^{n-2}r + \dots + r^{n-1}) > 0 \quad \leftarrow [-r, +\infty)$$

$$(a-r)(a+r)(a-r) > 0 \quad \frac{-r}{-1} \quad \frac{r}{+1} \Rightarrow a \cdot (-r, r) \cup (r, +\infty)$$

$f^{-1}(r) = r$, $f(n) = r_{n+k}$, $f^{-1}(r) = (k, 2)$, $f(e) = r \Rightarrow k+k = k-1$. (4)
 الف) $f(v) = r_{n-1, 2}$, $r_1 - 1, 2$. (5)

1) $f(f(n)) = 3(3n-1) - 1 = 9n - 4$

$$A: (ra, a) \quad , \quad f(n) = \frac{an}{n-1} \Rightarrow f^{-1}(n) \Rightarrow n = \frac{ay}{y-1} \Rightarrow ny - n = ay \quad (5) \quad (u)$$

$$-n = ay - ny \Rightarrow -n = y(a-n) \Rightarrow y = \frac{-n}{a-n} = \frac{n}{n-a} \quad \underline{\underline{(ra, a)}} \quad a = \frac{ra}{\frac{ra}{a}} \Rightarrow a = r$$

$$g = \{(u, r), (r, q), (q, w)\}, f = \{(u, w), (w, v), (v, r), (r, q)\}$$

$$\text{الف) } f \circ f^{-1} : \{(5,5), (7,7), (2,2), (9,9)\}$$

∴ $P^{-1} P = \{(r, r), (x, x), (v, v), (q, q)\}$

c.) $\log^{-1} = \{(2, 6), (6, 9)\}$

$$\Rightarrow g^{-1} \circ f_2 \{ (r, q), (v, r), (q, \wedge) \}$$

$$f = \{(r, r), (r, 1), (r, 0)\}, g = \{(r, 1), (r, r), (r, 0)\}, h = \{(1, r), (r, r), (r, 1), (0, 1)\}$$

$$\frac{h}{f \circ g^{-1}} = \left\{ \left(\frac{0}{1}, \frac{1}{r} \right), \left(r, \frac{r}{1} \right), \left(1, \frac{r}{r} \right) \right\} \Rightarrow \frac{h}{f \circ g^{-1}} = \left\{ \left(r, \frac{1}{r} \right), \left(1, \frac{1}{r} \right) \right\}$$

$y = \frac{r^2 n + 1}{n - r}$

$\frac{r n}{n} = \cos$

$n = r \cos \Rightarrow n = r \cos$

$n = r \cos \Rightarrow \frac{1}{r}$

$y = 1$

$$2. \frac{r_{y1}}{y-1} = 2r_{y2} - r_{y3} + (2r_1 - r_2 - r_3 - r_{y2})$$


$$\frac{r_{n-1}}{r-n} = y \Rightarrow y = \frac{r_{n-1}}{n-r} \Rightarrow \frac{1}{\mu}$$

$$f(n) = \begin{cases} 1 & n \geq 1 \\ r n - r & 1 \leq n < r \\ -r & n < 1 \end{cases} \Rightarrow n = (y - r) \Rightarrow n + r = y \Rightarrow y = \frac{n}{r} + r$$



(5)

$$f(n) = \begin{cases} n^r + r & n \geq 1 \\ r n - 1 & n \leq 0 \end{cases}$$



$$f^{-1}(n) = \begin{cases} n^{\frac{1}{r}} + r & n \geq r+1 \\ \frac{n+1}{r} & n \leq r \end{cases}$$

$$f^{-1}(n) = \begin{cases} y = \sqrt[r]{n-r} & n \geq r \\ \frac{n+1}{r} & n \leq r \end{cases}$$

$$D_f = (-\infty, 0] \cup [r, +\infty)$$

$$R_f = (-\infty, -1] \cup [r, +\infty)$$

$$D_{f^{-1}} = (-\infty, -1] \cup [r, +\infty)$$

$$R_{f^{-1}} = (-\infty, 0] \cup [1, +\infty)$$

$$f(n) = n^r - \frac{(n+1)^r}{n+r}, \quad f^{-1}(n) = \frac{an+b}{r n+d}, \quad f^{-1}(b) = ?$$

$$f(n) = n^r - \frac{n^r + r n^{r-1} - 1 - r n^{r-1} - r n}{n+r} = \frac{-r n - 1}{n+r}$$

$$f^{-1}(n) = y = \frac{-r y - 1}{y+r} \Rightarrow n y + r n = -r y - 1 \Rightarrow r n + 1 = -r y - n y \Rightarrow \frac{r n + 1}{-n - r} = y \Rightarrow y = \frac{r}{-r} = -1$$

$$y = \frac{-r n + r}{r n + r} \Rightarrow b = -r \Rightarrow f^{-1}(-r) = \frac{1-r}{r} = \omega$$

$$f(n) = \frac{n}{n^r + 1} \Rightarrow f^{-1}(n) = n = \frac{y}{y^r + 1} \Rightarrow n y^r + n = y \Rightarrow n y^r - y + n = 0$$

$$y = \frac{1 \pm \sqrt{1 - 4 n^r}}{2 n^r}$$

(5)