


(۱)



$$\Rightarrow r \sin \alpha \times \sin \alpha \times \Delta t \Rightarrow r \sin^2 \alpha \times \frac{1}{r} \times \Delta t \Rightarrow r^2 \leq 11 \Rightarrow r = \sqrt{11}$$

$$\Rightarrow r_1 + r_2 + r_3 + r_4 \leq 11 \Rightarrow 1 \times \sqrt{11} = 1 \times \sqrt{11} \Rightarrow \boxed{r = \sqrt{11}}$$

(۲)

$$\omega \times v \times \frac{1}{r} \times \sin A - v \times \frac{1}{r} \times \sin A = 1, v \omega \Rightarrow \frac{1}{r} \times \sin A (\omega v - v) \leq 1, v \omega$$

$$\Rightarrow \sin A (\omega) = \omega v \Rightarrow \sin A \leq \frac{1}{r}$$

$$\Rightarrow A = 10 - 6 \times 3$$

$$\Rightarrow A < 90 \Rightarrow \boxed{A = 30} \Rightarrow \boxed{\tan \alpha = \frac{\sqrt{10}}{r}}$$

(۳)


$$\frac{|\sin \alpha|}{|\cos \alpha|} \leq -\frac{1}{\cot \alpha} \Rightarrow \frac{\cos \alpha}{|\sin \alpha|} \leq -\cot \alpha \Rightarrow \text{در } \alpha \text{ ربع دوم}$$

$$\frac{1}{\sqrt{\cos^2 \alpha}} - \tan \alpha \leq \frac{1 + \sin \alpha}{|\cos \alpha|} \Rightarrow \boxed{\text{در } \alpha \text{ ربع دوم}}$$

$$\frac{1}{|\cos \alpha|} - \tan \alpha = \frac{1}{|\cos \alpha|} + \frac{\sin \alpha}{|\cos \alpha|}$$


(۴)

$$\tan\left(\frac{\pi}{2} - \alpha\right) = \cot(\alpha) = \cot(180 - B) = -\cot(B) = \boxed{-\frac{r}{v \omega}}$$



(۵)

$$\frac{r \cos(\pi v \omega - \pi r) - r \sin(180 - \pi r)}{\sin(180 + \pi r) - \cos(\pi v \omega + \pi r)} = \frac{-r \sin(\pi r) - r \sin(\pi r)}{-\sin(\pi r) - \sin(\pi r)} \leq \frac{-\omega}{-r} \Rightarrow \frac{\omega}{r} \Rightarrow \boxed{r, \omega}$$



$$\sin \alpha = \frac{\sqrt{d}}{r}$$

$$\tan \alpha = \frac{\sqrt{d}}{r}$$

$$\frac{\sin\left(\alpha + \frac{\pi}{2}\right) - \sin(\alpha - \pi)}{|\tan^2 \alpha - 1|} \leq \frac{\cos \alpha + \sin \alpha}{|\tan^2 \alpha - 1|} \leq \frac{\frac{r}{r} - \frac{\sqrt{d}}{r}}{\left(\left(\frac{\sqrt{d}}{r}\right)^2 - 1\right)}$$

$$= \frac{r - \sqrt{d}}{r} \leq \frac{r(\pi - \sqrt{d})}{r}$$

$$\tan \alpha = \frac{r \cos \alpha}{r \sin \alpha} = r \Rightarrow \tan \alpha + r = \frac{1}{r \sin \alpha} \quad (V)$$


$$\Rightarrow r+1 = \frac{1}{r \sin \alpha} \Rightarrow r \sin \alpha = \frac{1}{\sqrt{a}} \Rightarrow \boxed{\frac{-1}{\sqrt{a}}}$$

$$r m \sin \alpha + (m^2 - 1) y = r \Rightarrow y = \frac{r}{m^2 - 1} - \frac{r m}{m^2 - 1} x \quad (A)$$

$$\Rightarrow \frac{-r m}{m^2 - 1} = \tan \gamma \Rightarrow \frac{-r m}{m^2 - 1} = \sqrt{r} \Rightarrow -r m = \sqrt{r} m^2 - \sqrt{r}$$

$$\Rightarrow \sqrt{r} m^2 + r m - \sqrt{r} = 0 \Rightarrow \frac{\sqrt{a}}{|a|} = \boxed{\frac{r}{\sqrt{r}}}$$

$$\tan\left(\frac{\pi}{2} - \alpha\right) = \frac{1 - \tan \alpha}{1 + \tan \alpha} \xrightarrow{\substack{t = m \\ r = m}} \Rightarrow r - r \tan \alpha + m - m \tan \alpha = |r \tan \alpha - m - m \tan \alpha|$$



$$\Rightarrow r m \sin^2 \alpha - 1 \Rightarrow m = \frac{r \tan \alpha - 1}{r}$$

$$\Rightarrow -1 < \tan \alpha < 1 \Rightarrow -r < r \tan \alpha < r \Rightarrow -r < r \tan \alpha - 1 < r$$

$$\Rightarrow \boxed{-r < \frac{r \tan \alpha - 1}{r} < 1}$$

$$\begin{aligned} \tan \pi &= \tan(\pi - \gamma) = -\tan(\gamma) = -\sqrt{r} \\ \cos \pi &= \cos(\pi + \pi) = -\cos \pi = -\frac{\sqrt{r}}{r} \\ \tan \pi &= \tan(\pi - \gamma) = -\tan \gamma = -\sqrt{r} \\ \sin \pi &= \sin(\pi + \pi) = \sin \pi = \frac{\sqrt{r}}{r} \end{aligned} \quad \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} \begin{aligned} &(-\sqrt{r}) \left(-\frac{\sqrt{r}}{r}\right) + (-\sqrt{r}) \left(\frac{\sqrt{r}}{r}\right) \\ &= \boxed{0} \end{aligned} \quad (1)$$