

$$(4) A = \mu x - 1 = \frac{\mu}{\mu} - \frac{1}{\mu} x \rightarrow \frac{\omega}{\mu} x = \frac{\omega}{\mu} \rightarrow x = 1 \rightarrow y = 1$$

$$B = \frac{\mu}{\mu} - \frac{1}{\mu} x = \varepsilon x \rightarrow x = \omega \rightarrow y = -1$$

$$C = \varepsilon x = \mu x - 1 \rightarrow \mu x = \omega \rightarrow x = \frac{\omega}{\mu} \rightarrow y = \frac{\mu}{\mu}$$

$$M \left| \frac{1}{\mu} \right| \rightarrow AM = \sqrt{\frac{\varepsilon^4}{q} + \frac{1}{q}} = \frac{\omega \sqrt{\mu}}{\mu}$$

$$MBC = 1 \rightarrow y = x - 9$$

$$MAH = -1 \rightarrow y = -x + \mu \rightarrow H = x - 9 = -x + \mu \rightarrow x = \varepsilon \rightarrow y = \mu$$

$$AH = \sqrt{\mu^2 + \mu^2} = \mu \sqrt{2} \rightarrow \frac{AM}{AH} = \frac{\frac{\omega \sqrt{\mu}}{\mu}}{\mu \sqrt{2}} = \frac{\omega}{q}$$

$$(V) m = -\frac{1}{\mu} \rightarrow y = \frac{1}{\mu} x + C \xrightarrow{x^2} x^2 + \mu y - \mu C = 0$$

$$x = 0 \rightarrow \mu y = \mu C \rightarrow y = C \rightarrow (0, C)$$

$$y = 0 \rightarrow x = \mu C \rightarrow (\mu C, 0)$$

$$d = \frac{|1(0) + \mu(0) - \mu C|}{\sqrt{1 + \mu^2}} = \frac{|-\mu C|}{\sqrt{2}} = \sqrt{2} \rightarrow |\mu C| = \sqrt{2} \cdot \varepsilon$$

$$\mu C = 1 \rightarrow C = \omega \rightarrow \sqrt{(1-\omega)^2 + (\omega-\omega)^2} = \sqrt{1-2\omega}$$

$$(9) \xrightarrow{B(x_0)} y = \frac{1+x'_B}{\mu} \rightarrow x'_B = \mu y - 1$$

$$-1 = \frac{0+y'_B}{\mu} \rightarrow y'_B = -\varepsilon$$

$$(1) m = -1 \rightarrow y = x - (x-1) \rightarrow y = -x + \mu$$

$$M = -x + \mu = x + \omega \rightarrow x = -\mu \rightarrow x = -1 \rightarrow y = \varepsilon$$


$$M \left| \frac{1+4}{\mu} \right| = -1 \rightarrow a = -\mu$$

$$\left| \frac{\mu+b}{\mu} \right| = \varepsilon \rightarrow b = \mu$$

$$\mu b - a = 1 + \mu = 1 + \omega$$

$$\frac{d}{1} + x \frac{b}{1} \rho_2 - (m-x) \frac{3}{1} = 1 - \rho$$

$$\frac{b}{1} \rightarrow c, m$$



$$h = s - \frac{t}{1t} = m$$

$$m(m, 1) w$$