



$$A \begin{vmatrix} -2 \\ K \end{vmatrix}$$

$$B \begin{vmatrix} 2 \\ m \end{vmatrix}$$

$$A \begin{vmatrix} -2 \\ K \end{vmatrix}$$

$$B \begin{vmatrix} 2 \\ -2+K \end{vmatrix}$$

(1)

$$\text{if } M_{AB} = -\frac{1}{f} \Rightarrow \frac{m-k}{\epsilon + \frac{1}{f}} = -\frac{1}{f} \rightarrow m-k = -\frac{\epsilon + \frac{1}{f}}{f}$$

$$m = -\frac{\epsilon + \frac{1}{f}}{f}$$

$$\overline{AB} = \sqrt{(4)^2 + (-3)^2} = \sqrt{16+9} = \sqrt{25} = 5$$

5

$$A \begin{vmatrix} -1 \\ \epsilon \end{vmatrix}$$

$$B \begin{vmatrix} 3 \\ 1 \end{vmatrix}$$

$$C \begin{vmatrix} x \\ y \end{vmatrix}$$

$$D \begin{vmatrix} -1-x \\ y+3 \end{vmatrix}$$

P, P

(2)

$$M_{AB} = M(CD) \rightarrow \frac{3}{-\epsilon} = \frac{-3}{2x+1}$$

$$2x+1 = -\epsilon$$

$$x = \frac{-\epsilon - 1}{2}$$

$$M_{AB} \times M(CD) = -1 \rightarrow \frac{3}{-\epsilon} \times \frac{-1-y}{y+3} = -1 \rightarrow y = -1$$

$$C \begin{vmatrix} \frac{3}{f} \\ -1 \end{vmatrix}$$

$$\overline{AB} = \sqrt{16+9} = 5$$

$$\overline{BC} = \sqrt{\frac{9}{\epsilon^2} + \epsilon^2} = \frac{3}{\epsilon}$$

$$\rightarrow P = \left(\frac{3}{\epsilon}, \frac{3}{\epsilon} \right) = \left(\frac{3}{\epsilon}, \frac{3}{\epsilon} \right)$$

$$x^2 + (x^2 - 1)y = 3$$

(3)

15

$$\text{y. n. j.} \Rightarrow \tan \phi = \frac{1}{\sqrt{3}} \rightarrow \phi = 30^\circ$$

باجت (4)

$$\frac{-a}{b} \rightarrow \frac{-2m}{m^2-1} = \sqrt{3}$$

$$\sqrt{3}m^2 - \sqrt{3} = -2m$$

$$m = \frac{-2 \pm \sqrt{4 - 12}}{2\sqrt{3}} \rightarrow 1) \frac{-2}{\sqrt{3}}$$

$$\Delta = 4 - 12 = -8$$

$$\left| \frac{-2}{\sqrt{3}} - \frac{1}{\sqrt{3}} \right| = \frac{3}{\sqrt{3}} = \sqrt{3}$$

20

$$MBC = \frac{11-3}{\sqrt{10}} = \frac{8}{\sqrt{10}} = 2.5$$

(C) 1

$$y = 2x + b \xrightarrow{|x|} y = 2x - 3 \Rightarrow b = -3$$

(D)

$$y = 2x + 3 \rightarrow A \mid 4$$

BC is A line

$$\frac{|4 - 2 + 3|}{\sqrt{1+4}} = \frac{5}{\sqrt{5}} = \sqrt{5}$$

$$AB: y = -2x + 5 \xrightarrow{x=1} y = 3$$

$$BC: y = 2x - 3$$

$$\Rightarrow -2x + 5 = 2x - 3$$

$$B \mid 1$$


$$y = 1 \quad \boxed{x = 3} \leftarrow 11x = 33$$

(P)

$$AC: y = -2x - 1$$

$$\frac{|-1 - 2 - 1|}{\sqrt{5}} = \frac{4}{\sqrt{5}} = \frac{4\sqrt{5}}{5}$$

10



$$m = \frac{y}{x} = \frac{-1}{-1} = 1$$

14

$$y = -1x + b \xrightarrow{y=0} y = -1 \Rightarrow b = -1$$

$$x \rightarrow y = 1x - 1$$

$$y = -1x - 1$$

(D)

15

$$x = -1x - 1 \rightarrow x = -\frac{1}{2}$$

$$\frac{1}{\sqrt{2}}$$

$$\sqrt{\left(\frac{1}{\sqrt{2}}\right)^2 + \left(\frac{1}{\sqrt{2}}\right)^2} = \sqrt{\frac{1}{2} + \frac{1}{2}} = \sqrt{1} = 1$$

$$y = ax + 1$$

$$y = \frac{1}{a}x + \frac{a-1}{a} \quad (v) \quad 1$$

$$ay = ax + a - 1$$

$$\text{if } a = 1 \rightarrow y = x + 1 \rightarrow y = x$$

صفر ممکنه

$$\text{if } a = -1 \rightarrow y = -x + 1, y = -x + 2$$

$$\parallel \alpha \rightarrow \text{صفر ممکنه}$$

5

$$\text{if } a = 1 \rightarrow x + 1 = y, y = x \rightarrow y = x - 1, y = x$$

$$\text{فاصله دو خط موازی} = \frac{1}{\sqrt{1}} = \frac{\sqrt{2}}{2} \rightarrow 0^2 - \left(\frac{\sqrt{2}}{2}\right)^2 = \left(\frac{\sqrt{2}}{2}\right)^2$$

$$\frac{1}{\sqrt{2}} = \text{فاصله} \leftarrow \sqrt{2 \cdot 0 - 1} = \frac{1}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{1}{\sqrt{2}}$$

10

$$x - 3y = 1 \rightarrow 3y = x - 1 \rightarrow y = \frac{x}{3} - \frac{1}{3} \quad (5) \quad (1)$$

$$d = \frac{|1 \cdot 1 - 1 \cdot 1|}{\sqrt{1^2 + 9}} = \frac{0}{\sqrt{10}} = 0$$

$$MAC = -3 \rightarrow y = -2x + b \xrightarrow{b=2} y = -2x + 1$$


$$3y = x - 1 \rightarrow 3y = -9x + 24 \rightarrow 1 \cdot x = 3y \rightarrow \begin{matrix} y = 1 \\ x = 3 \end{matrix}$$

$$1/9 - 3/4 \cdot 2 = -2/1 \quad S = \frac{1}{2} \times 2 \cdot 1/1 = 1/1$$

$$\left| \frac{-1}{2} \right|$$

$$\left| \frac{-1}{3} \right|$$

$$\frac{b-a}{\left(-\frac{1}{2} / \frac{1}{3}\right)} = \sqrt{3} \quad \sqrt{\frac{1}{3^2} (b-a)^2} = \frac{\sqrt{3}}{3}$$



$$\sqrt{\frac{1}{3^2} + \frac{1}{1^2}} = \sqrt{\frac{1}{9} + 1} = \sqrt{\frac{10}{9}} = \frac{\sqrt{10}}{3}$$

$$2 \sqrt{\frac{10}{3^2}} = \frac{2\sqrt{10}}{3}$$

20

Subject: _____

Date: _____

$$\begin{aligned} & \begin{vmatrix} -r \\ -f \end{vmatrix} \\ & r = \sqrt{\mu^2 + f^2} = 0 \quad M_O A = M_B C \quad (1.) \end{aligned}$$

$$BC \rightarrow y = \frac{f}{\mu} x + b \xrightarrow{\mu} \mu y = f x + \mu b$$

$$r = \frac{|\mu y - f x + \mu b|}{\sqrt{f^2}} = 0 \quad |\mu b| = r_0 \quad b = \frac{r_0}{\mu} \checkmark$$

$$b = -\frac{r_0}{\mu} \quad \leftarrow$$

$$BC = y \Rightarrow \frac{f}{\mu} x + \frac{r_0}{\mu} \xrightarrow{\mu} \mu y - f x - r_0 = 0$$

$$AB \rightarrow y = -\frac{f}{\mu} x + b \quad \begin{vmatrix} -f \\ -\mu \end{vmatrix}$$

$$b = -\frac{r_0}{\mu} \quad \leftarrow$$

$$AB \rightarrow \begin{cases} \mu x + f y + r_0 \\ \mu y - f x - r_0 \end{cases}$$

$$\rightarrow \begin{aligned} x &= -\frac{-r_0 - 1 \dots}{-12 - 9} = -\sqrt{10} \\ y &= \frac{1 \dots - r_0}{-12 - 9} = -1 \rightarrow \mu y = \sqrt{10} \end{aligned}$$